

Chapter 5

PRO-APOTOTIC AND ANTI-APOPTOITIC EFFECTS OF TUMOR NECROSIS FACTOR IN TUMOR CELLS

Role of Nuclear Transcription Factor NF- κ B

Bharat B. Aggarwal and Yasunari Takada

Cytokine Research Section, Department of Experimental Therapeutics, The University of Texas, M.D. Anderson Cancer Center, Houston, TX

1. INTRODUCTION

An intricate balance between cell growth and cell death drives the proper growth, development, and function of most tissues¹. A vast amount of information has accumulated regarding the molecular mechanisms governing cell growth, but the mechanisms by which cells regulate their own death still remain a matter of great intrigue and have recently begun to acquire great importance. One known mechanism, apoptosis, or programmed cell death, is a physiological process believed to be responsible for the deletion of unwanted cells during organ and tissue development, tissue homeostasis and removal of self-reactive immune cells and pathologically induced tissue damage. Virus-infected cells are eliminated by the interaction with cytotoxic T-lymphocytes that kill the virus infected cells by inducing apoptosis^{2,3}. Cells that have DNA damage undergo apoptosis so as to eliminate cells that have accumulated genetic mutations and may become cancerous^{4,5}. In addition to being activated during development-related cell reduction, apoptosis can be triggered in many cell types by various stresses, including chemotherapeutic agents, cytokines, ionizing radiation, osmotic stress, and expression of viral proteins such as E1A⁶.

Extensive research within the last few years has revealed that cell death, whether at the single cell level, the tissue/organ level, or the organism level, is as important to life as cell survival. The critical role of apoptosis has been recognized in a wide variety of situations including immunomodulation, autoimmunity, sepsis, arthritis, inflammatory bowel disease, chronic heart failure, periodontal diseases, allograft rejection, neovascularization, obesity, tumorigenesis, meningitis, and parturition⁷.

NF- κ B is a ubiquitously expressed transcription factor that plays a pivotal role in expression of various inducible target genes that regulate apoptosis among several other vital functions⁸ it also controls, cell proliferation, differentiation, and immune and inflammatory responses. This

factor is a member of the Rel family of proteins, which bind to specific DNA sequences. In non-stimulated cells, the heterodimeric NF- κ B complexes are sequestered in the cytoplasm of most cell types by inhibitory proteins of the I κ B family (**Figure 1**)⁹.




Figure 1. Negative regulation of apoptosis by the NF- κ B-regulated gene products.

These inhibitors mask the NF- κ B nuclear localization domain and inhibit its DNA-binding activity. In response to a large variety of stimuli, the I κ B inhibitor is rapidly phosphorylated and degraded, thus allowing NF- κ B nuclear translocation, DNA binding to specific recognition sequences in promoters, and transcription of the target genes^{10,11}. Rel/NF- κ B transcription factors are induced in response to a large variety of stimuli and regulate a number of genes. The Rel/NF- κ B transcription factor family is comprised of several structurally related proteins that exist in organisms from insects to humans. The vertebrate family includes five cellular proteins: c-Rel, RelA, RelB, p50/p105, and p52/p100. These proteins can form homodimers or heterodimers giving diverse combinations of dimeric complexes that bind to DNA target sites, collectively called κ B sites, and directly regulate gene expression. The most common transcription factor of this family is called NF- κ B and consists of a p50/RelA heterodimer. The different Rel/NF- κ B proteins show distinct ability to form dimers, distinct preferences for different κ B sites, and distinct abilities to bind to I κ B inhibitor proteins¹². Thus, different Rel/NF- κ B complexes can be induced in different cell types and by distinct signals (**Figure 2**), can interact in distinct ways with other transcription factors and regulatory proteins, and can regulate the expression of distinct gene sets. Numerous kinases have been implicated in the activation of NF- κ B induced by different agents (**Figure 3**). Furthermore, the activation of NF- κ B is regulated both negatively and positively by other transcription factors and gene products (**Figure 4**).




Figure 2. Positive regulation of apoptosis by the NF- κ B-regulated gene products.




Figure 3. Regulation of NF- κ B activation by various protein kinases.




Figure 4. Regulation of NF- κ B activation by p65-binding proteins

Mechanism of apoptosis: Besides dying by necrosis, multicellular organisms can initiate a series of events that activate intracellular proteases and ultimately result in the destruction of the cell. These are collectively known as apoptosis. Apoptotic cells undergo an orderly series of biochemical or morphological events including cell shrinkage, mitochondrial breakdown, and nuclear DNA fragmentation¹³. The dying cell degrades into subcellular membrane-bound vesicles called apoptotic bodies, which are ultimately removed by phagocytosis. Apoptosis is a molecular suicide program characterized by cytoplasmic shrinkage, nuclear condensation, and DNA fragmentation into 180-base pair fragments¹⁴⁻¹⁷. It is a genetically regulated mechanism, and its deregulation can result in multistep carcinogenesis¹⁸⁻²⁰.

Apoptosis is brought about by activation of the family of proteins known as caspases (cysteiny, aspartate-specific proteases)^{21,22}. There are about 14 caspases involved in the process of apoptosis. Caspases are synthesized as pro-enzymes that are activated by proteolysis at two or three sites to remove an N-terminal peptide and divide the pro-enzyme into large and small subunits, which in some cases are joined by a linker domain. The mature caspase is a heterotetramer of two large and two small subunits^{23,24}. All caspases are activated by cleavage at a specific aspartate residue and act in a cascade. They are ultimately responsible for the proteolysis of the cellular substrates responsible for apoptosis.

Poly (ADP-ribose) polymerase (PARP) is the most well characterized substrate for several caspase in many cell systems. Intact PARP (116 kDa) is cleaved into two fragments (89 kDa and 24 kDa) during apoptosis^{25,26}. Cleavage of PARP is a valuable indicator of apoptosis, but its biological relevance is not known. Caspase-activated deoxyribonuclease (CAD) is a cytoplasmic endonuclease whose activation is thought to be responsible for generating the oligonucleosomal DNA fragments that are the hallmark of apoptosis²⁷.

DNA-dependent protein kinase (DNA-PK) is a DNA repair enzyme that is degraded during apoptosis by caspase-3²⁸. Degradation of DNA-PK will result in a decrease in the capacity of the cell to repair damage of nuclear DNA, thus facilitating the breakdown of DNA that is associated with apoptosis. Caspase-6 is responsible for degradation of lamin, which are the major structural components of the nuclear envelope²⁹. Cleavage of the cytoskeletal proteins fodrin³⁰, Gas 2³¹, and actin³² during apoptosis may induce cell shrinkage and membrane blebbing and alter cell signaling pathways. U1-70kDa, a small ribonucleosomal particle that functions in the splicing of mRNA transcripts, is cleaved during apoptosis (**Figure 1 and Figure 2**)³³. Caspases also cleave the initiation factors³⁴. This may inhibit translation during apoptosis. Caspases also cleave certain cell-signaling proteins, e.g., PKC- α and MEKK-1, which are rendered constitutively active and pro-apoptotic. In contrast, protein kinase B, which is involved in the anti-apoptotic pathway, is cleaved and inactivated by caspases³⁵.

A cell is induced to undergo apoptosis either by internal signals arising within the cells or external signals triggered by death activators that bind to receptors located at the cell surface. Internal signals initiate apoptosis in the mitochondria with the release of cytochrome c^{36,37}. The mitochondrial pathway is controlled by the Bcl-2 family of proteins³⁸. There are 15 members of the Bcl-2 protein family that share homology in at least one of three conserved domains (BH1–BH4) and these may either promote survival e.g., Bcl-2, Bcl-x_L or promote apoptosis, e.g., Bax, or Bak³⁹. The Bcl-2 family of proteins register both positive and negative stimuli and integrate them to determine whether the mitochondrial apoptotic pathway is turned on or off. Oncogenes encode mutated versions of the signaling proteins that control normal cell proliferation e.g., Ras signaling. Another, the Raf oncoprotein eventually initiates apoptosis when the cell receives an abnormal proliferative signal⁴⁰.

The apoptotic program can also be initiated by the action of extracellular messengers, termed death ligands. These bind to the cell surface receptors, termed death receptors, that activate intracellular signaling events that begin an apoptotic cascade^{41,42}. Death receptors belong to the TNF receptor superfamily that is characterized by a cysteine-rich extracellular ligand-binding domain⁴³. Death receptors contain a consensus module known as the death domain that is found in the intracellular portion of the molecule and is involved in transducing the apoptotic signal⁶. Fas and the TNF receptor are the two best-characterized death receptors, the cognate ligands for which are FasL and TNF, respectively.

Among all the known physiological inducers of apoptosis in mammalian cells, tumor necrosis factor (TNF) is perhaps the most potent and well studied. Many other members of the TNF superfamily also induce apoptosis, including LT (lymphotoxin), FasL (fibroblast-associated ligand), TRAIL (TNF-related apoptosis-inducing ligand), DR3L (for death receptor 3 ligand or also known as TWEAK for a weak homologue of TNF), THANK (TNF homologue that activates apoptosis, NF- κ B and JNK), and VEGI (vascular endothelial cell growth inhibitor)^{44,45}. Whether all these TNF family members induce apoptosis by the same mechanism as TNF is not known. Besides killer cytokines outlined above, apoptosis is also induced by various chemotherapeutic agents.

Within the last few years, a series of biochemical steps have been identified in the apoptotic pathway induced by cytokines and chemotherapeutic agents. For instance in TNF-induced apoptosis the TNF receptor is activated, which, through its cytoplasmic death domain, recruits a protein called TNF receptor-associated death domain (TRADD), which in turn sequentially recruits Fas-associated death domain (FADD) and FADD-like ICE (FLICE, also called caspase-8)⁴⁶⁻⁴⁸. The last activates caspase-9, which in turn activates caspase-3 (the executioner protease), resulting in apoptosis.

In contrast to cytokines, chemotherapeutic agents induce cellular apoptosis by inducing formation of mitochondrial transition pores, a rapid decrease in the mitochondrial transmembrane potential, and release of cytochrome c. The latter, in the presence of the protein Apaf-1, activates caspase-9, which then activates caspase-3. Several recent studies, however, have suggested that these two receptor-mediated and non-receptor-mediated pathways initiated by cytokines and chemotherapeutic agents, respectively, are not exclusive of each other and share similar steps.

Most agents that induce apoptosis also activate NF- κ B. Thus it is not too surprising that almost all cytokines of the TNF superfamily and chemotherapeutic agents activate NF- κ B. TNF-induced activation of NF- κ B (primarily consisting of p50 and p65 subunits) involves recruitment of TNF receptor-associated factor (TRAF)-2 by TRADD, which then binds to NIK. TRADD also binds to receptor-interacting protein (RIP). Either NIK or RIP then activate a kinase called I κ B kinase (IKK), which in turn leads to the phosphorylation, ubiquitination, and degradation of I κ B (the inhibitory subunit of NF- κ B), leading to NF- κ B activation⁴⁸. Some recent studies exclude NIK from a role in TNF-induced NF- κ B activation. How chemotherapeutic agents activate NF- κ B is not fully understood, but most likely it also involves phosphorylation, ubiquitination, and degradation of I κ B. How NF- κ B activation is linked with induction of apoptosis by TNF and chemotherapeutic agents is the subject of this review.

Anti-apoptotic effects of NF- κ B: Almost five years ago it was shown that TNF-induced apoptosis can be blocked by NF- κ B activation⁴⁹⁻⁵². Rel/NF- κ B transcription factors exercise their anti-apoptotic effects in a wide variety of cells to protect them from various apoptotic agents. They promote cell survival by inducing the transcription of anti-apoptotic genes (**Figure 1**). Activation of NF- κ B either upregulates the activity of anti-apoptotic genes or downregulates the activity of apoptotic genes. Inhibition of NF- κ B nuclear translocation enhances apoptotic killing by cytokines that belong to the TNF superfamily, ionizing radiation, overexpression of oncoproteins, chemotherapeutic agents, cytokines, phorbol esters, hyperoxia, hormones, and micro-organisms (**Table 1**, at the end of this chapter).

Some earlier studies showed that the oncogene v-rel from the avian retrovirus reticuloendotheliosis virus strain can block apoptosis⁵³ in chickens. Similarly, v-rel rendered chicken B-cells resistant to radiation-induced apoptosis⁵⁴. A large number of reports have demonstrated the anti-apoptotic effect of NF- κ B in a wide variety of cell types. The protective role of NF- κ B has now been shown in a large variety of cell types, including human breast carcinoma⁵⁰, T-cells^{51,55,56}, fibroblasts and macrophages⁴⁹, endothelial cells⁵⁷, EBV-infected lymphoblastoid cells⁵⁸, non-small lung cancer cells⁵⁹, glomerular mesangial cells⁶⁰, human ovarian cancer cells⁶¹,

human pancreatic cancer cell lines ⁶², Ewing sarcoma cells ⁶³, cardiomyocytes ⁶⁴, mouse embryos ⁶⁵, and HT1080 fibrosarcoma ⁵².

Treatment of RelA-deficient (the transcriptionally active subunit of NF- κ B) mouse fibroblasts and macrophages with TNF significantly reduced cell viability, whereas RelA^{+/+} cells were unaffected. In addition, reintroduction of RelA into RelA^{-/-} fibroblasts enhanced survival, demonstrating that Rel A is required for protection from TNF ⁴⁹. Another report showed that activation of the NF- κ B by TNF, ionizing radiation, or daunorubicin protects cells from apoptosis, whereas inhibition of NF- κ B enhanced apoptotic killing by these reagents but not by apoptotic stimuli that do not activate NF- κ B ⁵². Van Antwerp *et al.*, however, showed that the sensitivity and kinetics of TNF-induced apoptosis are enhanced in a number of cell types expressing a dominant-negative I κ B β (an inhibitor of NF- κ B) ⁵¹. Continued expression of v-Rel is necessary to maintain the viability of transformed lymphoid cells and enables primary spleen cells to escape apoptosis in culture ⁶⁶.

Liu *et al.* used the signaling proteins and showed that recruitment of FADD to the TNFR1 complex mediates apoptosis, that recruitment of RIP and TRAF2 mediate NF- κ B activation, and that activation of the latter protects cells against TNF-induced apoptosis ⁵⁰. Substoichiometric TFIID subunit TAFII105 is essential for activation of anti-apoptotic genes in response to TNF- α , serving as a transcriptional co-activator for NF- κ B ⁶⁷.

Adenovirus E1A protein has inhibited activation of NF- κ B and rendered cells more sensitive to TNF-induced apoptosis. This inhibition was brought about through suppression of I κ B kinase (IKK) activity and I κ B phosphorylation ⁶⁸. NF- κ B can attenuate TNF- α -induced apoptosis without de novo protein synthesis in the human pancreatic cancer cell lines MIA PaCa-2 and Capan-2. TNF- α -induced apoptosis was blocked by IL-1 β , a potent inducer of NF- κ B activation ⁶². These findings suggest that de novo protein synthesis is dispensible for anti-apoptotic effects of NF- κ B and support the possibility that NF- κ B exerts its anti-apoptotic action through protein-protein interaction.

The NF- κ B cascade is important in Bcl-x_L expression and for the anti-apoptotic effects of the CD28 receptor in primary human CD4⁺ lymphocytes ⁵⁶. HuT-78, a lymphoblastoid T-cell line with constitutive NF- κ B activity, contains elevated levels of Bcl-x_L protein and, similar to proliferating CD4⁺ T-cells, is resistant to apoptotic stimuli such as anti-Fas and TNF- α . In contrast, the same stimuli readily induced apoptosis in Jurkat cells without producing any detectable Bcl-x_L expression.

The quinone reductase inhibitors dicoumarol and menadione block SAPK/JNK and NF- κ B and thereby potentiate apoptosis ⁶⁹. Javelaud and Besancon have demonstrated that the repression of JNK activation by NF- κ B is involved in the anti-apoptotic effect of this transcription factor in TNF- α -treated Ewing sarcoma cells ⁶³. Also, NF- κ B exercises its anti-apoptotic effects through NF- κ B-inducing kinases (NIK). NIK induces PC12 cell differentiation and prevents apoptosis ⁷⁰. Cardiomyocytes utilize transcription factor NF- κ B to activate survival factors in the context of TNF- α stimulation. As locally increased levels of TNF- α have been detected in heart failure, NF- κ B activity is essential for cellular homeostasis in the heart

NF- κ B is required for TNF-mediated induction of the gene encoding human cIAP2. When overexpressed in mammalian cells, cIAP2 activates NF- κ B and suppresses TNF cytotoxicity. Both of these cIAP2 activities are blocked in vivo by coexpression of a dominant form of I κ B that is resistant to TNF-induced degradation⁵⁵. Functional coupling of NF- κ B and cIAP2 during the TNF response may provide signal amplification loop that promotes cell survival rather than death. The *IAP* genes function to protect the cell from undergoing apoptotic death in response to a variety of stimuli. The *IAP* genes *hIAP1*, *hIAP2*, and *XIAP* were found to be strongly upregulated upon treatment of endothelial cells with the inflammatory cytokines TNF- α , IL-1 α , and LPS, which in turn lead to activation of NF- κ B. This suggests that xiap represents one of the NF- κ B-regulated genes that counteracts the apoptotic signals elicited by TNF- α and thereby prevents endothelial cells from undergoing apoptosis during inflammation⁵⁷.

Treatment of WEHI 231 cells with N-tosyl-L-phenylalanine chloromethyl ketone, a protease inhibitor that prevents degradation of I κ B, or with low doses of pyrrolidine dithiocarbamate selectively inhibited NF- κ B activation and induced apoptosis⁷¹. Similarly, microinjection of WEHI 231 cells with either I κ B-GST protein or a c-Rel affinity-purified antibody induced apoptosis⁷¹.

Arlt *et al.* have shown that under certain conditions the resistance of pancreatic carcinoma cells to chemotherapy is due to their constitutive NF- κ B rather than the transient induction of NF- κ B by some anti-cancer drugs⁷². Exposure of normal keratinocytes to IFN- γ plus TPA produced a synergistic activation of NF- κ B. They acquired a resistance to UV-light-induced apoptosis that was dependent on NF- κ B because expression of a dominant negative form of I κ B overcame the resistance⁷³. There is enough evidence to suggest that activation and proper regulation of NF- κ B is essential for acquisition of an apoptotic-resistant phenotype for epidermal-derived keratinocytes. Kolenko *et al.* have demonstrated that inhibition of NF- κ B activity by cell permeable SN50 peptide in human T lymphocytes induces caspase-dependent apoptosis⁷⁴. Kawai *et al.* have shown that p53 is involved in NF- κ B inactivation and is required for x-ray-induced apoptosis in thymic lymphoma cells and normal thymocytes⁷⁵.

Oxidative stress induces apoptosis in human aortic endothelial cells through the downregulation of Bcl-2, translocation of Bax, and upregulation of p53, probably through NF- κ B activation. Oxidative stress may play an important role in endothelial apoptosis mediated by hypoxia, through the activation of NF- κ B⁷⁶. NF- κ B is a redox-sensitive transcription factor that is activated by oxidative insult, and NF- κ B activation can protect cells from apoptosis. When human alveolar epithelial (A549) cells were exposed to hyperoxia, NF- κ B was activated and within minutes was translocated to the nucleus⁷⁷. Reactive oxygen species could act synergistically with TNF α in causing cytotoxicity via inhibition of a cytoprotective branch of TNF α signaling pathways that starts with NF- κ B activation. Ginis *et al.* have demonstrated that H₂O₂ inhibited TNF α -induced accumulation of p65 in the nucleus, although it had no effect on degradation of I κ B in the cytoplasm⁷⁸.

It is known that adenovirus protein E1B blocks TNF-induced apoptosis, whereas E1A enhances TNF-induced apoptosis through unknown

mechanisms. Recent evidence indicates the effect of these proteins is mediated through modulation of NF- κ B activation⁶¹.

The growth arrest-specific 6 gene product (Gas6) is a growth and survival factor related to protein S. Gas6 induces a rapid and transient increase in nuclear NF- κ B binding activity coupled to transcription activation. This plays a central role in promoting survival in NIH 3T3 cells⁷⁹. MKK6 activates myocardial cell NF- κ B and inhibits apoptosis in a p38 mitogen-activated protein kinase dependent manner⁸⁰. Limb girdle muscular dystrophy type 2A results in decreased production of calpain 3. Calpain 3 is responsible for I κ B α turnover. Over expression of I κ B α results in sequestration of NF- κ B outside the nucleus. Myonuclear apoptosis occurred because of the downregulation of NF- κ B⁸¹.

The stimulation of the CD95- and TRAIL-resistant human pancreatic adenocarcinoma cell line Panc Tu1 with an agonistic anti-CD95 antibody or TRAIL activates of protein kinase C and NF- κ B. The activation of PKC operates directly in a death receptor dependent manner in PancTu1 cells and pancreatic tumor cells, protecting them from anti-CD95 and TRAIL-mediated apoptosis by preventing the loss of Δ and cytochrome c release as well as by induction of NF- κ B⁸². Pharmacologic or molecular inhibition of the NF- κ B pathway blocked cell survival in MCF-7 APO⁺ cells, while only molecular inhibition induced cytotoxicity in the APO⁻ cells⁴⁰. TGF- β protected gastric mucosal cells against apoptosis induced by serum depletion or sodium butyrate in a dose-dependent manner. This anti-apoptotic effect of TGF- β was blocked by pre-treatment with reagents that can potentially inhibit NF- κ B activation. This suggests that TGF- β plays an anti-apoptotic role in gastric mucosal cells via the NF- κ B-dependent pathway⁸³.

Mice deficient in the NF- κ B2 gene were challenged with the intracellular parasite *Toxoplasma gondii*. During the chronic phase of the infection, susceptibility of NF- κ B knockout mice to toxoplasmic encephalitis was associated with a reduced capacity of their splenocytes to produce IFN- γ associated with a loss of CD4⁺ and CD8⁺ T-cells. This loss of T-cells correlated with increased levels of apoptosis and with elevated expression of the pro-apoptotic molecule Fas by T-cells from infected NF- κ B knockout mice. This suggests a role of NF- κ B in maintenance of T-cell responses required for long-term resistance to *Toxoplasma gondii*⁸⁴.

How NF- κ B suppresses apoptosis? Although it is clear that NF- κ B activation plays a role in suppressing TNF-induced apoptosis, just how is only now beginning to emerge. Several genes that may play a role in blocking apoptosis and whose expression is regulated by NF- κ B have been identified, including cellular inhibitors of apoptosis (cIAP)-1 and cIAP-2, TRAF-1, and TRAF-2^{55,57,85}. cIAP-1, cIAP-2, and TRAF-1 are known to bind to TRAF-2 and TRAF-2 is required for NF- κ B activation. Thus, how these proteins block apoptosis is not clear. Other reports show that TNF induces manganese superoxide dismutase (SOD), whose expression is also regulated by NF- κ B, and the overexpression of SOD induces resistance to TNF-induced apoptosis⁸⁶. Also, altered SOD expression in HeLa cells after low dose γ -irradiation is responsible for NF- κ B-mediated cisplatin resistance⁸⁷. Insulin manifests its anti-apoptotic signaling through the activation of the NF- κ B-dependent survival genes encoding TRAF-2 and SOD⁸⁸. The TNF-

inducible zinc finger protein A20 is regulated by NF- κ B, and the role of this protein in induction of resistance to TNF-induced apoptosis has been demonstrated^{89,90}. The expression of a protein critical in the regulation of the cell cycle, cyclin D1, is also regulated by NF- κ B, and this activity may contribute to the cell growth and differentiation function assigned to NF- κ B^{91,92}.

The pro-survival Bcl-2 homolog Bfl-1/A1 is another gene whose transcription is regulated by NF- κ B and blocks TNF-induced apoptosis^{93,94}. There are other studies which show that Bcl-2 activates NF- κ B through the degradation of the inhibitor I κ B⁹⁵. Crawford *et al.* have demonstrated that Bcl-2 overexpression protects photooxidative stress-induced apoptosis of photoreceptor cells through NF- κ B preservation. It has been known that the Ras/PI-3K/Akt pathway plays a critical role in cell survival. It now appears that this pathway is linked to the activation of IKK, the kinase needed for I κ B phosphorylation and NF- κ B activation. Akt may also play a cytoprotective role through activation of NF- κ B^{96,97}. An NF- κ B-independent cytoprotective pathway has also been described. The NF- κ B activation induced by overexpression of TRAF2 was found to be insufficient to protect cells from apoptosis induced by TNF and cycloheximide together, thus indicating an essential role for additional components in the cytoprotective response⁹⁸.

While NF- κ B activation blocks apoptosis, it seems that activation of apoptosis also blocks NF- κ B activation, suggesting a feedback loop. For instance, endothelial cells undergo apoptosis when deprived of growth factors. The surviving viable cells exhibit increased activity of NF- κ B, whereas apoptotic cells show caspase-mediated cleavage of the NF- κ B p65/RelA subunit, resulting in loss of carboxy-terminal transactivation domains and a transcriptionally inactive p65 molecule, which itself acts as a dominant-negative inhibitor of NF- κ B, promoting apoptosis. In contrast an uncleavable, caspase-resistant p65 protects the cells from apoptosis. The generation of a dominant-negative fragment of p65 during apoptosis may be an efficient pro-apoptotic feedback mechanism between caspase activation and NF- κ B inactivation⁹⁹. Similarly apoptosis has been shown to promote a caspase-induced amino-terminal truncation of I κ B that functions as a stable inhibitor of NF- κ B¹⁰⁰, thus further enhancing apoptosis. And Fas, another member of the TNF receptor family, was found to induce caspase-3-mediated proteolysis of both p50 and p65 subunits of NF- κ B in T Jurkat cells, thus sensitizing the cells to apoptosis¹⁰¹.

Pro-apoptotic activity of NF- κ B: The decision of life or death in response to an inducing signal within a cell is dependent upon a delicate balance of positive and negative influences. While there are several reports that NF- κ B activation protects cells from undergoing apoptosis induced by TNF or chemotherapeutic agents, there are also reports suggesting that NF- κ B activation mediates apoptosis in response to a variety of inducers in a number of cell types. For instance, in murine clonal osteoblasts NF- κ B activation mediated TNF-induced apoptosis¹⁰². The suppression of growth of CD34⁺ myeloid cells by TNF also correlated with NF- κ B activation¹⁰³. Apart from this, Fas activates NF- κ B and induces apoptosis in T-cell lines by signaling pathways distinct from those induced by TNF α ¹⁰⁴. Human

melanoma cells are protected against UV-induced apoptosis through downregulation of NF- κ B activity and Fas expression¹⁰⁵. Oxidative stress induced apoptosis in human aortic endothelial cells through the downregulation of Bcl-2, translocation of Bax, and upregulation of p53 probably takes place through NF- κ B activation. Oxidative stress may play an important role in endothelial apoptosis mediated by hypoxia, through the activation of NF- κ B⁷⁶. That the activation of NF- κ B is rather required for apoptosis has also been shown for other inducers such as H₂O₂^{106,107}. Similarly, H₂O₂-induced apoptosis was not suppressed by hyperoxia-induced NF- κ B activation⁷⁷. In pancreatic islets, A20 inhibited both apoptosis and NF- κ B activation induced by cytokines, suggesting that NF- κ B may actually mediate apoptosis¹⁰⁸. Apoptosis in HL-60 cells induced by chemotherapeutic agents such as etoposide or 1-beta-D-arabinofuranosylcytosine was also found to require NF- κ B activation, inasmuch as suppression of NF- κ B by PDTC also blocked apoptosis¹⁰⁹.

Recently, Stark *et al.* demonstrated that aspirin induces cell death by an active apoptotic process that involves nuclear translocation of NF- κ B preceding cell death¹¹⁰. *Helicobacter pylori* induces NF- κ B-mediated apoptosis in chronic gastritis¹¹¹. The apoptosis induced by alphavirus was also found to require the activation of NF- κ B, since the thiol agents and Bcl-2 blocked both activities¹¹². During adenoviral infection, NF- κ B mediates apoptosis through transcriptional activation of Fas¹¹³. Apoptosis in Ca⁺⁺ reperfusion injury of cultured astrocytes was also found to be mediated through NF- κ B activation¹¹⁴. The cell death-promoting role of NF- κ B has also been demonstrated in focal cerebral malaria¹¹⁵, as it has for induction of apoptosis by double-stranded-RNA-dependent protein-kinase (PKR)¹¹⁶. Lin *et al.* showed that NF- κ B can be pro-apoptotic or anti-apoptotic depending on the timing of modulating NF- κ B activity relative to the death stimulus¹¹⁷. How NF- κ B may mediate apoptosis is not clear, but the role of p53 and c-myc induction through NF- κ B has been demonstrated¹¹⁸. In addition, NF- κ B is required for the anti-CD3-mediated apoptosis of double-positive thymocytes through a pathway that involves the regulation of the anti-apoptotic gene Bcl-x_L¹¹⁹. c-myc has also been implicated in survival of certain cells such as hepatocytes¹²⁰. These observations suggest that NF- κ B activation not only negatively, but also positively regulates apoptosis. This idea has been further strengthened by studies on NMRI mice, Wistar rats and WI-38 fibroblasts in which aging induced a strong and consistent increase in the nuclear binding activity of NF- κ B¹²¹.

We recently showed that doxorubicin and its structural analogues WP631 and WP744, activate NF- κ B, and this activation is essential for apoptosis in myeloid (KBM-5) and lymphoid (Jurkat) cells (138). Because the anthracycline analogue (WP744), most active as a cytotoxic agent, was also most active in inducing NF- κ B activation and the latter preceded the cytotoxic effects, suggests that NF- κ B activation may mediate cytotoxicity. Second, receptor-interacting protein-deficient cells, which did not respond to doxorubicin-induced NF- κ B activation, were also protected from the cytotoxic effects of all the three anthracyclines. Third, suppression of NF- κ B activation by pyrrolidine dithiocarbamate, also suppressed the cytotoxic effects of anthracyclines. Fourth, suppression of NF- κ B activation by

NEMO-binding domain peptide, also suppressed the cytotoxic effects of the drug. Overall our results clearly demonstrated that NF- κ B activation and I κ B degradation are early events activated by doxorubicin and its analogues and that they play a critical pro-apoptotic role.

Evidence that apoptosis is unaffected by NF- κ B: There are increasing reports that NF- κ B activation plays little or no role in apoptosis. For instance, Cai *et al.* showed that overexpression of I κ B β , an inhibitor of NF- κ B, in human breast carcinoma MCF7 cells inhibits NF- κ B activation but not TNF-induced apoptosis. Similarly, in endothelial cells A20 inhibited NF- κ B activation without enhancing TNF-induced apoptosis¹²². LPS- and IL-1-induced prolongation in survival of endothelial cells did not require NF- κ B activation¹²³. The pro- and anti-apoptotic role of NF- κ B appears to be determined more by the nature of the death stimulus than by the origin of the tissue¹¹³. Bone morphogenetic protein (BMP)-2 and -4 inhibited TNF-mediated apoptosis by inhibiting caspase-8 activation in C2C12 cells, a pluripotent mesenchymal cell line that has potential to differentiate into osteoblasts depending on BMP stimulation. The BMP/Smad signaling pathway can inhibit TNF-mediated apoptosis independently of the pro-survival activity of NF- κ B. This suggests that BMPs not only stimulate osteoblast differentiation but also promote cell survival during the induction of bone formation, offering new insight into the biological functions of BMPs¹²⁴. There are proteins that associate with cytokine receptors such as SODD (for silencer of death domain)¹²⁵, sentris¹²⁶, and c-FLIP¹²⁷, that can also negatively regulate apoptosis, again independently of NF- κ B.

The redox-sensitive transcription factor Ref-1 plays a critical role in the survival of endothelial cells in response to hypoxia and cytokines including TNF α . Upregulation of Ref-1 promotes endothelial cell survival in response to hypoxia and TNF through NF- κ B-independent and NF- κ B-dependent signaling cascades¹²⁸. It has been observed in human non-small-cell lung carcinoma that apoptosis induced by topoisomerase poisons, e.g. Etoposide, is not mediated by NF- κ B but can be manipulated by proteasome inhibitors¹²⁹. Why NF- κ B plays a role in apoptosis induced by some agents and not others is not clear but suggests that the apoptotic pathway varies from one inducer to another and also perhaps from one cell type to another.

Conclusion: It is clear that apoptosis is regulated by mitochondria-dependent and -independent pathways involving a series of proteins that preexist in the cells. Most agents that induce apoptosis, also activate NF- κ B and the latter suppresses apoptosis in most cases. While it may appear paradoxical that the same agent could perform both functions, in reality it is not. The same stress that induces cells to die provokes a self-defense response in the cell. How NF- κ B plays an anti-apoptotic role in some cells, pro-apoptotic in others and no role in some requires further understanding. It is possible that activation of NF- κ B alone is not sufficient to regulate apoptosis and that other transcription factors are involved (141). Most NF- κ B-regulated genes (such as cyclooxygenase-2) play critical roles in inflammation, suggesting that inflammation can also negatively regulate apoptosis.

Abbreviations used: NF- κ B, nuclear factor- κ B; TNF, tumor necrosis factor; I κ B, inhibitor of NF- κ B; TRADD, TNF receptor-associated death

domain; NIK, NF- κ B-inducing kinase; TRAF2, TNF receptor-associated factor 2; SOD, superoxide dismutase; RIP, receptor interacting proteins; SODD, silencer of death domain; FADD, Fas-associated death domain; FLICE, FADD-like ICE; c-FLIP, cellular FLICE inhibitory protein; LT, lymphotoxin; FasL, fibroblast associated ligand; TRAIL, TNF-related apoptosis-inducing ligand; DR3L, death receptor 3 ligand; TWEAK, weak homologue of TNF; THANK, TNF homologue that activates apoptosis, NF- κ B and JNK; JNK, c-jun N-terminal kinase; VEGI, vascular endothelial cell growth inhibitor; cIAP, cellular inhibitors of apoptosis; PKR, double-stranded-RNA-dependent protein kinase; MEKK, mitogen-activated protein kinase/extracellular signal-regulated kinase kinase

Table 1. Anti-apoptotic activity of NF- κ B

Apoptosis Inducing Agent	Cell Type	Reference
TNF	Rel A ^{-/-} fibroblasts and macrophages	49
TNF	MCF-7	50
TNF	HEF, Jurkat, T24	51
TNF- α , radiation, daunorubicin	HT1080	52
TNF	Jurkat	55
TNF	CD4 ⁺ T lymphocytes	56
TNF	Endothelial cells	57
TNF	EBV infected lymphoblastoid cells	58
TNF	A549, MCF-7	59
TNF	Glomerular mesangial cells	60
α -radiation	(SK-OV-3.ipl) cells	61
TNF	MIAPaCa-2, Capan-2	62
TNF	Ewing sarcoma cells	63
TNF	Cardiomyocytes	64
TNF, IL-1	Mouse embryos	65
v-Rel inducers	HeLa cells, spleen cells	66
TNF	293	67
TNF	SK-OV-3.ipl	68
TNF	Human pulmonary macrophages	69
NIK suppression	PC12	70
TPCK, PDTC (NF- κ B blockers)	WEHI 231	71
TPA and IFN- γ	Keratinocytes	73
SN50 (NF- κ B blocker)	T Lymphocytes	74
x-ray irradiation	Lymphoma cells, Thymocytes	75
Hyperoxia	A549	77
TNF- α and ROI	Brain capillary endothelial cells	78
Gas 6 suppression	NIH 3T3	79
Anisomycin	Myocardial cells	80
Calpain 3 deficiency	Myogenic satellite cells	81
Anti-CD95	Panc Tu1	82
Serum depletion, sodium butyrate	GSM 06	83
<i>Toxoplasma gondii</i>	T-cells	84
Insulin	CHP overexpressing insulin receptor	88
TNF	Prostate carcinoma cells	130
TGF- β , serum withdrawal, anoikis, TNF- α	Mv1Lu and MDCK	131
Growth factor deprivation	Hematopoietic cells	132
v-Rel	Spleen cells, fibroblasts, C4-1	133

Apoptosis Inducing Agent	Cell Type	Reference
TRAIL	Renal Cell carcinoma	134
Hyperoxia, TNF- α	Lung epithelial cells	135
TNF	Endothelial cells	136

MCF-7, human breast carcinoma; Panc Tu1, human pancreatic adenocarcinoma; A549, nonsmall cell lung cancer; SK-OV-3.ipl, human ovarian cancer cell line was generated from ascites developed in *nu/nu* mouse by administering an intraperitoneal injection of SK-OV-3, a human ovarian carcinoma cell line; MIAPaCa-2 and Capan-2, human pancreatic cancer cell lines; HT1080, fibrosarcoma; Mv1Lu and MDCK, epithelial cells; C4-1 and WEHI 231, B-cells; PC12, rat adrenal pheochromocytoma; GSM o6, gastric mucosal cell line.

Table 2. Pro-Apoptotic Activity of NF- κ B

Inducing Agent	Cell Type	Reference
Oxidative stress	Aortic endothelial cells	76
TNF α , HTLV-1	Osteoblast cell line	102
Tax/TNF α		
TNF α	Myeloid leukemic cell lines	103
Fas/TNF α	CEM-C7	104
UV light	Human melanoma	105
H ₂ O ₂	Jurkat, CEM C7, Oligodendrocytes	106, 107
Etoposide	HL-60 and thymocytes	109
Aspirin	Colon cancer cells	110
<i>Helicobacter pylori</i>	Gastric epithelial cells	111
Sindbis-virus induction	AT-3	112, 117
Adenovirus	Hepatocytes	113
RO1	Astrocytes	114
Focal cerebral ischemia	Neurons (Mice Ischemic model)	115
PKR	BSC-40, 3T3	116
Kainic acid	Rat striatum	118
α -CD3	Thymocytes from mI κ B mice	119
Constitutive enhanced by etoposide	Immature Rat thymocytes	137
Doxorubicin	KBM-5, SH-SY5Y, IMR32	138, 139
Mullerian Inhibiting substance	T47D, MDA-MB-231	140

Jurkat, CEM-C7, human T-cells; HL-60, human promyelocytic leukemia; KBM-5, human myeloid; SH-SY5Y, IMR32, N-type neuroblastoma cells; T47D, MDA-MB-231, human breast; BSC-40, African green monkey kidney cells; AT-3, prostrate carcinoma cell line; PKR, double-stranded-RNA-dependent protein kinase.

REFERENCES

1. Jacobson M. D., Weil M., Raff M. C.: Programmed cell death in animal development. *Cell*. 88:347-354, 1997
2. Darmon A. J., Nicholson D. W., Bleackley R. C.: Activation of the apoptotic protease CPP32 by cytotoxic T-cell-derived granzyme B. *Nature*. 377:446-448, 1995
3. Shibata S., Kyuwa S., Lee S. K., Toyoda Y., Goto N.: Apoptosis induced in mouse hepatitis virus-infected cells by a virus-specific CD8⁺ cytotoxic T-lymphocyte clone. *J Virol*. 68:7540-7545, 1994
4. Jaattela M.: Escaping cell death: survival proteins in cancer. *Exp Cell Res*. 248:30-43, 1999
5. Stambolic V., Mak T. W., Woodgett J. R.: Modulation of cellular apoptotic potential: contributions to oncogenesis. *Oncogene*. 18:6094-6103, 1999
6. Nagata S.: Apoptosis by death factor. *Cell*. 88:355-365, 1997

7. Aggarwal B. B., Vilcek J. Tumore Necrosis Factor: Structure, Function and Mechanism of Action. New York: Marcel Dekker; 1992
8. Ghosh S., May M. J., Kopp E. B.: NF-kappa B and Rel proteins: evolutionarily conserved mediators of immune responses. *Annu Rev Immunol.* 16:225-260, 1998
9. Beg A. A., Baldwin A. S., Jr.: The I kappa B proteins: multifunctional regulators of Rel/NF-kappa B transcription factors. *Genes Dev.* 7:2064-2070, 1993
10. Karin M.: How NF-kappaB is activated: the role of the IkappaB kinase (IKK) complex. *Oncogene.* 18:6867-6874, 1999
11. Pahl H. L.: Activators and target genes of Rel/NF-kappaB transcription factors. *Oncogene.* 18:6853-6866, 1999
12. Chen F. E., Ghosh G.: Regulation of DNA binding by Rel/NF-kappaB transcription factors: structural views. *Oncogene.* 18:6845-6852, 1999
13. Arends M. J., Wyllie A. H.: Apoptosis: mechanisms and roles in pathology. *Int Rev Exp Pathol.* 32:223-254, 1991
14. Kerr J. F., Wyllie A. H., Currie A. R.: Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br J Cancer.* 26:239-257, 1972
15. Clarke P. G., Clarke S.: Historic apoptosis. *Nature.* 378:230, 1995
16. Cohen J. J.: Apoptosis. *Immunol Today.* 14:126-130, 1993
17. Chinnaiyan A. M., Dixit V. M.: The cell-death machine. *Curr Biol.* 6:555-562, 1996
18. McDonnell T. J., Korsmeyer S. J.: Progression from lymphoid hyperplasia to high-grade malignant lymphoma in mice transgenic for the t(14; 18). *Nature.* 349:254-256, 1991
19. Thompson C. B.: Apoptosis in the pathogenesis and treatment of disease. *Science.* 267:1456-1462, 1995
20. Gilmore T. D., Koedood M., Piffat K. A., White D. W.: Rel/NF-kappaB/IkappaB proteins and cancer. *Oncogene.* 13:1367-1378, 1996
21. Cohen G. M.: Caspases: the executioners of apoptosis. *Biochem J.* 326 (Pt 1):1-16, 1997
22. Green D. R.: Apoptotic pathways: paper wraps stone blunts scissors. *Cell.* 102:1-4, 2000
23. Bhardwaj A, Aggarwal BB. Receptor-mediated choreography of life and death. *J Clin Immunol.* 23:317-32, 2003.
24. Cerretti D. P., Kozlosky C. J., Mosley B., Nelson N., Van Ness K., Greenstreet T. A., March C. J., Kronheim S. R., Druck T., Cannizzaro L. A., et al.: Molecular cloning of the interleukin-1 beta converting enzyme. *Science.* 256:97-100, 1992
25. Kaufmann S. H.: Induction of endonucleolytic DNA cleavage in human acute myelogenous leukemia cells by etoposide, camptothecin, and other cytotoxic anticancer drugs: a cautionary note. *Cancer Res.* 49:5870-5878, 1989
26. Kaufmann S. H., Desnoyers S., Ottaviano Y., Davidson N. E., Poirier G. G.: Specific proteolytic cleavage of poly(ADP-ribose) polymerase: an early marker of chemotherapy-induced apoptosis. *Cancer Res.* 53:3976-3985, 1993
27. Enari M., Sakahira H., Yokoyama H., Okawa K., Iwamatsu A., Nagata S.: A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD. *Nature.* 391:43-50, 1998
28. Song Q., Lees-Miller S. P., Kumar S., Zhang Z., Chan D. W., Smith G. C., Jackson S. P., Alnemri E. S., Litwack G., Khanna K. K., Lavin M. F.: DNA-dependent protein kinase catalytic subunit: a target for an ICE-like protease in apoptosis. *Embo J.* 15:3238-3246, 1996
29. Takahashi A., Alnemri E. S., Lazebnik Y. A., Fernandes-Alnemri T., Litwack G., Moir R. D., Goldman R. D., Poirier G. G., Kaufmann S. H., Earnshaw W. C.: Cleavage of lamin A by Mch2 alpha but not CPP32: multiple interleukin 1 beta-converting enzyme-related proteases with distinct substrate recognition properties are active in apoptosis. *Proc Natl Acad Sci U S A.* 93:8395-8400, 1996
30. Martin S. J., O'Brien G. A., Nishioka W. K., McGahon A. J., Mahboubi A., Saido T. C., Green D. R.: Proteolysis of fodrin (non-erythroid spectrin) during apoptosis. *J Biol Chem.* 270:6425-6428, 1995
31. Brancolini C., Benedetti M., Schneider C.: Microfilament reorganization during apoptosis: the role of Gas2, a possible substrate for ICE-like proteases. *Embo J.* 14:5179-5190, 1995

32. Mashima T., Naito M., Fujita N., Noguchi K., Tsuruo T.: Identification of actin as a substrate of ICE and an ICE-like protease and involvement of an ICE-like protease but not ICE in VP-16-induced U937 apoptosis. *Biochem Biophys Res Commun.* 217:1185-1192, 1995
33. Casciola-Rosen L., Nicholson D. W., Chong T., Rowan K. R., Thornberry N. A., Miller D. K., Rosen A.: Apopain/CPP32 cleaves proteins that are essential for cellular repair: a fundamental principle of apoptotic death. *J Exp Med.* 183:1957-1964, 1996
34. Antoku K., Liu Z., Johnson D. E.: Inhibition of caspase proteases by CrmA enhances the resistance of human leukemic cells to multiple chemotherapeutic agents. *Leukemia.* 11:1665-1672, 1997
35. Widmann C., Gibson S., Johnson G. L.: Caspase-dependent cleavage of signaling proteins during apoptosis. A turn-off mechanism for anti-apoptotic signals. *J Biol Chem.* 273:7141-7147, 1998
36. Green D. R., Reed J. C.: Mitochondria and apoptosis. *Science.* 281:1309-1312, 1998
37. Susin S. A., Zamzami N., Kroemer G.: Mitochondria as regulators of apoptosis: doubt no more. *Biochim Biophys Acta.* 1366:151-165, 1998
38. Adams J. M., Cory S.: The Bcl-2 protein family: arbiters of cell survival. *Science.* 281:1322-1326, 1998
39. Gross A., McDonnell J. M., Korsmeyer S. J.: BCL-2 family members and the mitochondria in apoptosis. *Genes Dev.* 13:1899-1911, 1999
40. Weldon C. B., Burrow M. E., Rolfe K. W., Clayton J. L., Jaffe B. M., Beckman B. S.: NF-kappa B-mediated chemoresistance in breast cancer cells. *Surgery.* 130:143-150, 2001
41. Sheikh M. S., Fornace A. J., Jr.: Death and decoy receptors and p53-mediated apoptosis. *Leukemia.* 14:1509-1513, 2000
42. Ashkenazi A., Dixit V. M.: Death receptors: signaling and modulation. *Science.* 281:1305-1308, 1998
43. Smith C. A., Farrah T., Goodwin R. G.: The TNF receptor superfamily of cellular and viral proteins: activation, costimulation, and death. *Cell.* 76:959-962, 1994
44. Mukhopadhyay A., Ni J., Zhai Y., Yu G. L., Aggarwal B. B.: Identification and characterization of a novel cytokine, THANK, a TNF homologue that activates apoptosis, nuclear factor-kappaB, and c-Jun NH2-terminal kinase. *J Biol Chem.* 274:15978-15981, 1999
45. Haridas V., Shrivastava A., Su J., Yu G. L., Ni J., Liu D., Chen S. F., Ni Y., Ruben S. M., Gentz R., Aggarwal B. B.: VEGI, a new member of the TNF family activates nuclear factor-kappa B and c-Jun N-terminal kinase and modulates cell growth. *Oncogene.* 18:6496-6504, 1999
46. Rath P. C., Aggarwal B. B.: TNF-induced signaling in apoptosis. *J Clin Immunol.* 19:350-364, 1999
47. Darnay B. G., Aggarwal B. B.: Signal transduction by tumour necrosis factor and tumour necrosis factor related ligands and their receptors. *Ann Rheum Dis.* 58 Suppl 1:12-113, 1999
48. Wallach D., Varfolomeev E. E., Malinin N. L., Goltsev Y. V., Kovalenko A. V., Boldin M. P.: Tumor necrosis factor receptor and Fas signaling mechanisms. *Annu Rev Immunol.* 17:331-367, 1999
49. Beg A. A., Baltimore D.: An essential role for NF-kappaB in preventing TNF-alpha-induced cell death. *Science.* 274:782-784, 1996
50. Liu Z. G., Hsu H., Goeddel D. V., Karin M.: Dissection of TNF receptor 1 effector functions: JNK activation is not linked to apoptosis while NF-kappaB activation prevents cell death. *Cell.* 87:565-576, 1996
51. Van Antwerp D. J., Martin S. J., Kafri T., Green D. R., Verma I. M.: Suppression of TNF-alpha-induced apoptosis by NF-kappaB. *Science.* 274:787-789, 1996
52. Wang C. Y., Mayo M. W., Baldwin A. S., Jr.: TNF- and cancer therapy-induced apoptosis: potentiation by inhibition of NF-kappaB. *Science.* 274:784-787, 1996
53. Gilmore T. D.: Multiple mutations contribute to the oncogenicity of the retroviral oncoprotein v-Rel. *Oncogene.* 18:6925-6937, 1999
54. Neiman P. E., Thomas S. J., Loring G.: Induction of apoptosis during normal and neoplastic B-cell development in the bursa of Fabricius. *Proc Natl Acad Sci U S A.* 88:5857-5861, 1991

55. Chu Z. L., McKinsey T. A., Liu L., Gentry J. J., Malim M. H., Ballard D. W.: Suppression of tumor necrosis factor-induced cell death by inhibitor of apoptosis c-IAP2 is under NF-kappaB control. *Proc Natl Acad Sci U S A*. 94:10057-10062, 1997
56. Khoshnan A., Tindell C., Laux I., Bae D., Bennett B., Nel A. E.: The NF-kappa B cascade is important in Bcl-xL expression and for the anti-apoptotic effects of the CD28 receptor in primary human CD4+ lymphocytes. *J Immunol*. 165:1743-1754, 2000
57. Stehlik C., de Martin R., Kumabashiri I., Schmid J. A., Binder B. R., Lipp J.: Nuclear factor (NF)-kappaB-regulated X-chromosome-linked iap gene expression protects endothelial cells from tumor necrosis factor alpha-induced apoptosis. *J Exp Med*. 188:211-216, 1998
58. Asso-Bonnet M., Feuillard J., Ferreira V., Bissieres P., Tarantino N., Korner M., Raphael M.: Relationship between IkappaBalpha constitutive expression, TNFalpha synthesis, and apoptosis in EBV-infected lymphoblastoid cells. *Oncogene*. 17:1607-1615, 1998
59. Lee K. Y., Chang W., Qiu D., Kao P. N., Rosen G. D.: PG490 (triptolide) cooperates with tumor necrosis factor-alpha to induce apoptosis in tumor cells. *J Biol Chem*. 274:13451-13455, 1999
60. Sugiyama H., Savill J. S., Kitamura M., Zhao L., Stylianou E.: Selective sensitization to tumor necrosis factor-alpha-induced apoptosis by blockade of NF-kappaB in primary glomerular mesangial cells. *J Biol Chem*. 274:19532-19537, 1999
61. Shao R., Karunakaran D., Zhou B. P., Li K., Lo S. S., Deng J., Chiao P., Hung M. C.: Inhibition of nuclear factor-kappaB activity is involved in E1A-mediated sensitization of radiation-induced apoptosis. *J Biol Chem*. 272:32739-32742, 1997
62. Kajino S., Suganuma M., Teranishi F., Takahashi N., Tetsuka T., Ohara H., Itoh M., Okamoto T.: Evidence that de novo protein synthesis is dispensable for anti-apoptotic effects of NF-kappaB. *Oncogene*. 19:2233-2239, 2000
63. Javelaud D., Besancon F.: NF-kappa B activation results in rapid inactivation of JNK in TNF alpha-treated Ewing sarcoma cells: a mechanism for the anti-apoptotic effect of NF-kappa B. *Oncogene*. 20:4365-4372, 2001
64. Bergmann M. W., Loser P., Dietz R., von Harsdorf R.: Effect of NF-kappa B Inhibition on TNF-alpha-induced apoptosis and downstream pathways in cardiomyocytes. *J Mol Cell Cardiol*. 33:1223-1232, 2001
65. Li Z. W., Chu W., Hu Y., Delhase M., Deerinck T., Ellisman M., Johnson R., Karin M.: The IKKbeta subunit of IkappaB kinase (IKK) is essential for nuclear factor kappaB activation and prevention of apoptosis. *J Exp Med*. 189:1839-1845, 1999
66. Zong W. X., Farrell M., Bash J., Gelinas C.: v-Rel prevents apoptosis in transformed lymphoid cells and blocks TNFalpha-induced cell death. *Oncogene*. 15:971-980, 1997
67. Yamit-Hezi A., Dikstein R.: TAFII105 mediates activation of anti-apoptotic genes by NF-kappaB. *Embo J*. 17:5161-5169, 1998
68. Shao R., Hu M. C., Zhou B. P., Lin S. Y., Chiao P. J., von Lindern R. H., Spohn B., Hung M. C.: E1A sensitizes cells to tumor necrosis factor-induced apoptosis through inhibition of IkappaB kinases and nuclear factor kappaB activities. *J Biol Chem*. 274:21495-21498, 1999
69. Cross J. V., Deak J. C., Rich E. A., Qian Y., Lewis M., Parrott L. A., Mochida K., Gustafson D., Vande Pol S., Templeton D. J.: Quinone reductase inhibitors block SAPK/JNK and NFkappaB pathways and potentiate apoptosis. *J Biol Chem*. 274:31150-31154, 1999
70. Foehr E. D., Lin X., O'Mahony A., Geleziunas R., Bradshaw R. A., Greene W. C.: NF-kappa B signaling promotes both cell survival and neurite process formation in nerve growth factor-stimulated PC12 cells. *J Neurosci*. 20:7556-7563, 2000
71. Wu M., Lee H., Bellas R. E., Schauer S. L., Arsura M., Katz D., FitzGerald M. J., Rothstein T. L., Sherr D. H., Sonenshein G. E.: Inhibition of NF-kappaB/Rel induces apoptosis of murine B cells. *Embo J*. 15:4682-4690, 1996
72. Arlt A., Vorndamm J., Breitenbroich M., Folsch U. R., Kalthoff H., Schmidt W. E., Schafer H.: Inhibition of NF-kappaB sensitizes human pancreatic carcinoma cells to apoptosis induced by etoposide (VP16) or doxorubicin. *Oncogene*. 20:859-868, 2001

73. Qin J. Z., Chaturvedi V., Denning M. F., Choubey D., Diaz M. O., Nickoloff B. J.: Role of NF-kappaB in the apoptotic-resistant phenotype of keratinocytes. *J Biol Chem.* 274:37957-37964, 1999
74. Kolenko V., Bloom T., Rayman P., Bukowski R., Hsi E., Finke J.: Inhibition of NF-kappa B activity in human T lymphocytes induces caspase-dependent apoptosis without detectable activation of caspase-1 and -3. *J Immunol.* 163:590-598, 1999
75. Kawai H., Yamada Y., Tatsuka M., Niwa O., Yamamoto K., Suzuki F.: Down-regulation of nuclear factor kappaB is required for p53-dependent apoptosis in X-ray-irradiated mouse lymphoma cells and thymocytes. *Cancer Res.* 59:6038-6041, 1999
76. Aoki M., Nata T., Morishita R., Matsushita H., Nakagami H., Yamamoto K., Yamazaki K., Nakabayashi M., Ogihara T., Kaneda Y.: Endothelial apoptosis induced by oxidative stress through activation of NF-kappaB: antiapoptotic effect of antioxidant agents on endothelial cells. *Hypertension.* 38:48-55, 2001
77. Li Y., Zhang W., Mantell L. L., Kazzaz J. A., Fein A. M., Horowitz S.: Nuclear factor-kappaB is activated by hyperoxia but does not protect from cell death. *J Biol Chem.* 272:20646-20649, 1997
78. Ginis I., Hallenbeck J. M., Liu J., Spatz M., Jaiswal R., Shohami E.: Tumor necrosis factor and reactive oxygen species cooperative cytotoxicity is mediated via inhibition of NF-kappaB. *Mol Med.* 6:1028-1041, 2000
79. Demarchi F., Verardo R., Varnum B., Brancolini C., Schneider C.: Gas6 anti-apoptotic signaling requires NF-kappa B activation. *J Biol Chem.* 276:31738-31744, 2001
80. Zechner D., Craig R., Hanford D. S., McDonough P. M., Sabbadini R. A., Glembotski C. C.: MKK6 activates myocardial cell NF-kappaB and inhibits apoptosis in a p38 mitogen-activated protein kinase-dependent manner. *J Biol Chem.* 273:8232-8239, 1998
81. Baghdiguian S., Martin M., Richard I., Pons F., Astier C., Bourg N., Hay R. T., Chemaly R., Halaby G., Loiselet J., Anderson L. V., Lopez de Munain A., Fardeau M., Mangeat P., Beckmann J. S., Lefranc G.: Calpain 3 deficiency is associated with myonuclear apoptosis and profound perturbation of the IkappaB alpha/NF-kappaB pathway in limb-girdle muscular dystrophy type 2A. *Nat Med.* 5:503-511, 1999
82. Trauzold A., Wermann H., Arlt A., Schutze S., Schafer H., Oestern S., Roder C., Ungefroren H., Lampe E., Heinrich M., Walczak H., Kalthoff H.: CD95 and TRAIL receptor-mediated activation of protein kinase C and NF-kappaB contributes to apoptosis resistance in ductal pancreatic adenocarcinoma cells. *Oncogene.* 20:4258-4269, 2001
83. Kanai M., Konda Y., Nakajima T., Izumi Y., Takeuchi T., Chiba T.: TGF-alpha inhibits apoptosis of murine gastric pit cells through an NF-kappaB-dependent pathway. *Gastroenterology.* 121:56-67, 2001
84. Caamano J., Tato C., Cai G., Villegas E. N., Speirs K., Craig L., Alexander J., Hunter C. A.: Identification of a role for NF-kappa B2 in the regulation of apoptosis and in maintenance of T cell-mediated immunity to *Toxoplasma gondii*. *J Immunol.* 165:5720-5728, 2000
85. Wang C. Y., Mayo M. W., Korneluk R. G., Goeddel D. V., Baldwin A. S., Jr.: NF-kappaB antiapoptosis: induction of TRAF1 and TRAF2 and c-IAP1 and c-IAP2 to suppress caspase-8 activation. *Science.* 281:1680-1683, 1998
86. Manna S. K., Zhang H. J., Yan T., Oberley L. W., Aggarwal B. B.: Overexpression of manganese superoxide dismutase suppresses tumor necrosis factor-induced apoptosis and activation of nuclear transcription factor-kappaB and activated protein-1. *J Biol Chem.* 273:13245-13254, 1998
87. Eichholtz-Wirth H., Sagan D.: IkappaB/NF-kappaB mediated cisplatin resistance in HeLa cells after low-dose gamma-irradiation is associated with altered SODD expression. *Apoptosis.* 5:255-263, 2000
88. Bertrand F., Desbois-Mouthon C., Cadoret A., Prunier C., Robin H., Capeau J., Atfi A., Cherqui G.: Insulin antiapoptotic signaling involves insulin activation of the nuclear factor kappaB-dependent survival genes encoding tumor necrosis factor receptor-associated factor 2 and manganese-superoxide dismutase. *J Biol Chem.* 274:30596-30602, 1999
89. Krikos A., Laherty C. D., Dixit V. M.: Transcriptional activation of the tumor necrosis factor alpha-inducible zinc finger protein, A20, is mediated by kappa B elements. *J Biol Chem.* 267:17971-17976, 1992

90. Opipari A. W., Jr., Hu H. M., Yabkowitz R., Dixit V. M.: The A20 zinc finger protein protects cells from tumor necrosis factor cytotoxicity. *J Biol Chem.* 267:12424-12427, 1992
91. Hinz M., Krappmann D., Eichten A., Heder A., Scheidereit C., Strauss M.: NF-kappaB function in growth control: regulation of cyclin D1 expression and G0/G1-to-S-phase transition. *Mol Cell Biol.* 19:2690-2698, 1999
92. Guttridge D. C., Albanese C., Reuther J. Y., Pestell R. G., Baldwin A. S., Jr.: NF-kappaB controls cell growth and differentiation through transcriptional regulation of cyclin D1. *Mol Cell Biol.* 19:5785-5799, 1999
93. Wang C. Y., Cusack J. C., Jr., Liu R., Baldwin A. S., Jr.: Control of inducible chemoresistance: enhanced anti-tumor therapy through increased apoptosis by inhibition of NF-kappaB. *Nat Med.* 5:412-417, 1999
94. Zong W. X., Edelstein L. C., Chen C., Bash J., Gelinas C.: The prosurvival Bcl-2 homolog Bfl-1/A1 is a direct transcriptional target of NF-kappaB that blocks TNFalpha-induced apoptosis. *Genes Dev.* 13:382-387, 1999
95. de Moissac D., Mustapha S., Greenberg A. H., Kirshenbaum L. A.: Bcl-2 activates the transcription factor NFkappaB through the degradation of the cytoplasmic inhibitor IkappaBalpha. *J Biol Chem.* 273:23946-23951, 1998
96. Romashkova J. A., Makarov S. S.: NF-kappaB is a target of AKT in anti-apoptotic PDGF signaling. *Nature.* 401:86-90, 1999
97. Yang C. H., Murti A., Pfeffer S. R., Kim J. G., Donner D. B., Pfeffer L. M.: Interferon alpha /beta promotes cell survival by activating nuclear factor kappa B through phosphatidylinositol 3-kinase and Akt. *J Biol Chem.* 276:13756-13761, 2001
98. Natoli G., Costanzo A., Guido F., Moretti F., Bernardo A., Burgio V. L., Agresti C., Levrero M.: Nuclear factor kB-independent cytoprotective pathways originating at tumor necrosis factor receptor-associated factor 2. *J Biol Chem.* 273:31262-31272, 1998
99. Levkau B., Scatena M., Giachelli C. M., Ross R., Raines E. W.: Apoptosis overrides survival signals through a caspase-mediated dominant-negative NF-kappa B loop. *Nat Cell Biol.* 1:227-233, 1999
100. Reuther J. Y., Baldwin A. S., Jr.: Apoptosis promotes a caspase-induced amino-terminal truncation of IkappaBalpha that functions as a stable inhibitor of NF-kappaB. *J Biol Chem.* 274:20664-20670, 1999
101. Ravi R., Bedi A., Fuchs E. J.: CD95 (Fas)-induced caspase-mediated proteolysis of NF-kappaB. *Cancer Res.* 58:882-886, 1998
102. Kitajima I., Soejima Y., Takasaki I., Beppu H., Tokioka T., Maruyama I.: Ceramide-induced nuclear translocation of NF-kappa B is a potential mediator of the apoptotic response to TNF-alpha in murine clonal osteoblasts. *Bone.* 19:263-270, 1996
103. Hu X., Tang M., Fisher A. B., Olashaw N., Zuckerman K. S.: TNF-alpha-induced growth suppression of CD34+ myeloid leukemic cell lines signals through TNF receptor type I and is associated with NF-kappa B activation. *J Immunol.* 163:3106-3115, 1999
104. Packham G., Lahti J. M., Fee B. E., Gawn J. M., Coustan-Smith E., Campana D., Douglas I., Kidd V. J., Ghosh S., Cleveland J. L.: Fas activates NF-kB and induces apoptosis in T-cell lines by signaling pathways distinct from those induced by TNF-a. *Cell Death Differ.* 4:130-139, 1997
105. Ivanov V. N., Ronai Z.: p38 protects human melanoma cells from UV-induced apoptosis through down-regulation of NF-kappaB activity and Fas expression. *Oncogene.* 19:3003-3012, 2000
106. Dumont A., Hehner S. P., Hofmann T. G., Ueffing M., Droge W., Schmitz M. L.: Hydrogen peroxide-induced apoptosis is CD95-independent, requires the release of mitochondria-derived reactive oxygen species and the activation of NF-kappaB. *Oncogene.* 18:747-757, 1999
107. Vollgraf U., Wegner M., Richter-Landsberg C.: Activation of AP-1 and nuclear factor-kappaB transcription factors is involved in hydrogen peroxide-induced apoptotic cell death of oligodendrocytes. *J Neurochem.* 73:2501-2509, 1999
108. Grey S. T., Arvelo M. B., Hasenkamp W., Bach F. H., Ferran C.: A20 inhibits cytokine-induced apoptosis and nuclear factor kappaB-dependent gene activation in islets. *J Exp Med.* 190:1135-1146, 1999

109. Bessho R., Matsubara K., Kubota M., Kuwakado K., Hirota H., Wakazono Y., Lin Y. W., Okuda A., Kawai M., Nishikomori R., et al.: Pyrrolidine dithiocarbamate, a potent inhibitor of nuclear factor kappa B (NF-kappa B) activation, prevents apoptosis in human promyelocytic leukemia HL-60 cells and thymocytes. *Biochem Pharmacol.* 48:1883-1889, 1994
110. Stark L. A., Din F. V., Zwacka R. M., Dunlop M. G.: Aspirin-induced activation of the NF-kappaB signaling pathway: a novel mechanism for aspirin-mediated apoptosis in colon cancer cells. *Faseb J.* 15:1273-1275, 2001
111. Gupta R. A., Polk D. B., Krishna U., Israel D. A., Yan F., DuBois R. N., Peek R. M., Jr.: Activation of peroxisome proliferator-activated receptor gamma suppresses nuclear factor kappa B-mediated apoptosis induced by Helicobacter pylori in gastric epithelial cells. *J Biol Chem.* 276:31059-31066, 2001
112. Lin K. I., Lee S. H., Narayanan R., Baraban J. M., Hardwick J. M., Ratan R. R.: Thiol agents and Bcl-2 identify an alphavirus-induced apoptotic pathway that requires activation of the transcription factor NF-kappa B. *J Cell Biol.* 131:1149-1161, 1995
113. Kuhnel F., Zender L., Paul Y., Tietze M. K., Trautwein C., Manns M., Kubicka S.: NFkappaB mediates apoptosis through transcriptional activation of Fas (CD95) in adenoviral hepatitis. *J Biol Chem.* 275:6421-6427, 2000
114. Takuma K., Lee E., Kidawara M., Mori K., Kimura Y., Baba A., Matsuda T.: Apoptosis in Ca2+ reperfusion injury of cultured astrocytes: roles of reactive oxygen species and NF-kappaB activation. *Eur J Neurosci.* 11:4204-4212, 1999
115. Schneider A., Martin-Villalba A., Weih F., Vogel J., Wirth T., Schwaninger M.: NF-kappaB is activated and promotes cell death in focal cerebral ischemia. *Nat Med.* 5:554-559, 1999
116. Gil J., Alcamí J., Esteban M.: Induction of apoptosis by double-stranded-RNA-dependent protein kinase (PKR) involves the alpha subunit of eukaryotic translation initiation factor 2 and NF-kappaB. *Mol Cell Biol.* 19:4653-4663, 1999
117. Lin K. I., DiDonato J. A., Hoffmann A., Hardwick J. M., Ratan R. R.: Suppression of steady-state, but not stimulus-induced NF-kappaB activity inhibits alphavirus-induced apoptosis. *J Cell Biol.* 141:1479-1487, 1998
118. Nakai M., Qin Z. H., Chen J. F., Wang Y., Chase T. N.: Kainic acid-induced apoptosis in rat striatum is associated with nuclear factor-kappaB activation. *J Neurochem.* 74:647-658, 2000
119. Hettmann T., DiDonato J., Karin M., Leiden J. M.: An essential role for nuclear factor kappaB in promoting double positive thymocyte apoptosis. *J Exp Med.* 189:145-158, 1999
120. Bellas R. E., Sonenshein G. E.: Nuclear factor kappaB cooperates with c-Myc in promoting murine hepatocyte survival in a manner independent of p53 tumor suppressor function. *Cell Growth Differ.* 10:287-294, 1999
121. Helenius M., Hanninen M., Lehtinen S. K., Salminen A.: Changes associated with aging and replicative senescence in the regulation of transcription factor nuclear factor-kappa B. *Biochem J.* 318 (Pt 2):603-608, 1996
122. Ferran C., Stroka D. M., Badrichani A. Z., Cooper J. T., Wrighton C. J., Soares M., Grey S. T., Bach F. H.: A20 inhibits NF-kappaB activation in endothelial cells without sensitizing to tumor necrosis factor-mediated apoptosis. *Blood.* 91:2249-2258, 1998
123. Zen K., Karsan A., Stempien-Otero A., Yee E., Tupper J., Li X., Eunson T., Kay M. A., Wilson C. B., Winn R. K., Harlan J. M.: NF-kappaB activation is required for human endothelial survival during exposure to tumor necrosis factor-alpha but not to interleukin-1beta or lipopolysaccharide. *J Biol Chem.* 274:28808-28815, 1999
124. Chen S., Guttridge D. C., Tang E., Shi S., Guan K., Wang C. Y.: Suppression of tumor necrosis factor-mediated apoptosis by nuclear factor kappaB-independent bone morphogenetic protein/Smad signaling. *J Biol Chem.* 276:39259-39263, 2001
125. Jiang Y., Woronicz J. D., Liu W., Goeddel D. V.: Prevention of constitutive TNF receptor 1 signaling by silencer of death domains. *Science.* 283:543-546, 1999
126. Okura T., Gong L., Kamitani T., Wada T., Okura I., Wei C. F., Chang H. M., Yeh E. T.: Protection against Fas/APO-1- and tumor necrosis factor-mediated cell death by a novel protein, sentrin. *J Immunol.* 157:4277-4281, 1996

127. Scaffidi C., Schmitz I., Krammer P. H., Peter M. E.: The role of c-FLIP in modulation of CD95-induced apoptosis. *J Biol Chem.* 274:1541-1548, 1999
128. Hall J. L., Wang X., Van A., Zhao Y., Gibbons G. H.: Overexpression of Ref-1 inhibits hypoxia and tumor necrosis factor-induced endothelial cell apoptosis through nuclear factor-kappaB-independent and -dependent pathways. *Circ Res.* 88:1247-1253, 2001
129. Tabata M., Tabata R., Grabowski D. R., Bukowski R. M., Ganapathi M. K., Ganapathi R.: Roles of NF-kappaB and 26 S proteasome in apoptotic cell death induced by topoisomerase I and II poisons in human nonsmall cell lung carcinoma. *J Biol Chem.* 276:8029-8036, 2001
130. Herrmann J. L., Beham A. W., Sarkiss M., Chiao P. J., Rands M. T., Bruckheimer E. M., Brisbay S., McDonnell T. J.: Bcl-2 suppresses apoptosis resulting from disruption of the NF-kappa B survival pathway. *Exp Cell Res.* 237:101-109, 1997
131. Lallemand F., Mazars A., Prunier C., Bertrand F., Kornprost M., Gallea S., Roman-Roman S., Cherqui G., Atfi A.: Smad7 inhibits the survival nuclear factor kappaB and potentiates apoptosis in epithelial cells. *Oncogene.* 20:879-884, 2001
132. Besancon F., Atfi A., Gespach C., Cayre Y. E., Bourgeade M. F.: Evidence for a role of NF-kappaB in the survival of hematopoietic cells mediated by interleukin 3 and the oncogenic TEL/platelet-derived growth factor receptor beta fusion protein. *Proc Natl Acad Sci USA.* 95:8081-8086, 1998
133. You M., Ku P. T., Hrdlickova R., Bose H. R., Jr.: ch-IAP1, a member of the inhibitor-of-apoptosis protein family, is a mediator of the antiapoptotic activity of the v-Rel oncoprotein. *Mol Cell Biol.* 17:7328-7341, 1997
134. Oya M., Ohtsubo M., Takayanagi A., Tachibana M., Shimizu N., Murai M.: Constitutive activation of nuclear factor-kappaB prevents TRAIL-induced apoptosis in renal cancer cells. *Oncogene.* 20:3888-3896, 2001
135. Franek W. R., Horowitz S., Stansberry L., Kazzaz J. A., Koo H. C., Li Y., Arita Y., Davis J. M., Mantell A. S., Scott W., Mantell L. L.: Hyperoxia inhibits oxidant-induced apoptosis in lung epithelial cells. *J Biol Chem.* 276:569-575, 2001
136. Hofer-Warbinek R., Schmid J. A., Stehlik C., Binder B. R., Lipp J., de Martin R.: Activation of NF-kappa B by XIAP, the X chromosome-linked inhibitor of apoptosis, in endothelial cells involves TAK1. *J Biol Chem.* 275:22064-22068, 2000
137. Slater A. F., Kimland M., Jiang S. A., Orrenius S.: Constitutive nuclear NF kappa B/rel DNA-binding activity of rat thymocytes is increased by stimuli that promote apoptosis, but not inhibited by pyrrolidine dithiocarbamate. *Biochem J.* 312 (Pt 3):833-838, 1995
138. Ashikawa K., Shishodia S., Fokt I., Priebe W., Aggarwal B. B.: Evidence that activation of nuclear factor-kappaB is essential for the cytotoxic effects of doxorubicin and its analogues. *Biochem Pharmacol.* 67:353-364, 2004
139. Bian X., McAllister-Lucas L. M., Shao F., Schumacher K. R., Feng Z., Porter A. G., Castle V. P., Opipari A. W., Jr.: NF-kappa B activation mediates doxorubicin-induced cell death in N-type neuroblastoma cells. *J Biol Chem.* 276:48921-48929, 2001
140. Segev D. L., Ha T. U., Tran T. T., Kenneally M., Harkin P., Jung M., MacLaughlin D. T., Donahoe P. K., Maheswaran S.: Mullerian inhibiting substance inhibits breast cancer cell growth through an NFkappa B-mediated pathway. *J Biol Chem.* 275:28371-28379, 2000
141. Aggarwal BB. Signalling pathways of the TNF superfamily: a double-edged sword *Nat Rev Immunol.* 2003 Sep; 3(9):745-56.